

**Online-Only Abstract: Population-based burden of bloodstream infections in Finland****Diagnosis, management and outcome of *Candida* endocarditis**

A. Lefort<sup>1</sup>, L. Chartier<sup>2</sup>, B. Sendid<sup>3</sup>, M. Wolff<sup>4</sup>, J.-L. Mainardi<sup>5,6,7</sup>, I. Podglajen<sup>5</sup>, M. Desnos-Ollivier<sup>8,9</sup>, A. Fontanet<sup>2,10</sup>, S. Bretagne<sup>8,9,11</sup>, O. Lortholary<sup>1,8,9</sup>, for the French Mycosis Study Group

1) Université Paris Descartes, AP-HP, Hôpital Necker-Enfants malades, Service des Maladies Infectieuses et Tropicales, Centre d'Infectiologie Necker-Pasteur, 2) Institut Pasteur, Unité d'Epidémiologie des Maladies Emergentes, Paris, 3) Université Nord de France, INSERM U995, CHRU, Université Lille II, Lille, 4) Université Paris Diderot, AP-HP, Hôpital Bichat, Service de Réanimation Médicale, 5) Université Paris Descartes, AP-HP, Hôpital Européen Georges Pompidou, Service de Microbiologie Clinique, 6) Université Pierre et Marie Curie, UMR S 872, 7) INSERM, U872, LRMA, Equipe 12 du Centre de Recherche des Cordeliers, 8) Institut Pasteur, Unité de Mycologie Moléculaire, Centre National de Référence Mycologie et Antifongiques, 9) CNRS URA3012, 10) Conservatoire National des Arts et Métiers, Chaire Santé et Développement, Paris and 11) Université Paris 12, AP-HP, Hôpital Henri Mondor, Laboratoire de Mycologie-Parasitologie, Créteil, France

**Original Submission:** 2 October 2011; **Revised Submission:** 3 December 2011; **Accepted:** 3 January 2012

Editor: D. Raoult

**Article published online:** 7 January 2012

*Clin Microbiol Infect* 2012; **18**: E99–E109

10.1111/j.1469-0691.2012.03764.x

**Abstract**

Limited data exist on *Candida* endocarditis (CE) outcome in the era of new antifungals. As early diagnosis of CE remains difficult, non-culture-based tools need to be evaluated. Through the French prospective MYCENDO study (2005–2007), the overall characteristics and risk factors for death from CE were analysed. The contribution of antigen detection (mannan/anti-mannan antibodies and (1,3)- $\beta$ -D-glucans) and molecular tools was evaluated. Among 30 CE cases, 19 were caused by non-*albicans* species. Sixteen patients (53%) had a predisposing cardiac disease, which was a valvular prosthesis in ten (33%). Nine patients (30%) were intravenous drug users; none of them had right-sided CE. Among the 21 patients who were not intravenous drug users, 18 (86%) had healthcare-associated CE. Initial therapy consisted of a combination of antifungals in 12 of 30 patients (40%). Thirteen patients (43%) underwent valve replacement. The median follow-up was 1 year after discharge from hospital (range, 5 months to 4 years) and hospital mortality was 37%. On univariate analysis, patients aged  $\geq 60$  years had a higher mortality risk (OR 11, 95% CI 1.2–103.9;  $p$  0.024), whereas intravenous drug use was associated with a lower risk of death (OR 0.12, 95% CI 0.02–0.7;  $p$  0.03). Among 18 patients screened for both serum mannan/anti-mannan antibodies and (1,3)- $\beta$ -D-glucans, all had a positive result with at least one of either test at CE diagnosis. Real-time PCR was performed on blood (SeptiFast) in 12 of 18, and this confirmed the blood culture results. In conclusion, CE prognosis remains poor, with a better outcome among younger patients and intravenous drug users. Detection of serum antigens and molecular tools may contribute to earlier CE diagnosis.